



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DIOPHANTINE ANALYSIS.

134. Proposed by O. E. GLENN, Ph. D., Springfield, Mo.

How many sets of solutions has the congruence $\lambda + \mu + \nu + \xi \equiv 0 \pmod{p-1}$ p being a prime number; the order of λ, μ, ν, ξ being disregarded.

GEOMETRY.

284. Proposed by JOHN JAMES QUINN, Ph. D., Warren, Pa.

a) Suppose that two radii R and r , whose center is the origin, revolve with uniform angular velocities 3θ and θ , respectively. What is the equation of the locus of P , the projection parallel to the X axis of the extremity of the radius r on the radius R produced if necessary.

b) Apply this curve to the trisection of an angle.

c) Suppose the ratio of their velocities is $n\theta:\theta$. Show how we can effect the multisection of an angle.

285. Proposed by G. E. BROCKWAY, Nashua, N. H.

Prove without the aid of the circle, that if the bisectors of the angles of a triangle be drawn, the greatest bisector falls on the least side.

286. Proposed by S. F. NORRIS, Baltimore City College, Baltimore, Md.

On the sides of a given triangle measure off equal distances from the extremities of the base, and at these points erect perpendiculars to the sides. Find the locus of the point of intersection of these perpendiculars. Solve by methods of analytic geometry.

287. Proposed by G. W. GREENWOOD, M. A., McKendree College, Lebanon, Ill.

Show that the points whose abscissae are 0, $a_1/\sqrt{3}$, and $-a_1/\sqrt{3}$ are points of inflexion on the locus $x^2y - a^2x + a^2y = 0$.

MISCELLANEOUS.

157. Proposed by H. L. ORCHARD, M. A., B. Sc. (Unsolved problem in the Educational Times, London.)

An inelastic rod 9 feet long is placed with its upper end upon a rough vertical plane and its lower end on a smooth horizontal plane, and so that it makes an angle of 45° with each plane. It is now released, and strikes against a smooth sphere of 1 foot diameter placed in contact with the two planes. Determine the subsequent motion.

UNSOLVED PROBLEMS.

NOTE. The following problems still remain unsolved (in our columns).

Group Theory, 8. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

In a chess tournament between eight players, there are seven rounds, the